

NALOXONE PROVIDERS

Find Statewide Distribution Sites

www.nvopioidresponse.org

Harm Reduction Agencies

Change Point

445 Ralston Street, Reno, NV 89503
775-997-7519

Trac-B Exchange

6114 W Charleston Blvd., Las Vegas, NV 89146
702-840-6693

Treatment Providers

www.behavioralhealthnv.org

The average cash pay price range
for NARCAN Nasal Spray in

Nevada is \$75-\$145

NARCAN Nasal Spray is covered under
NV Medicaid

For immediate support for behavioral
health crisis services call or text 988

Opioid Safety

and how to use naloxone



This project was supported in whole or in part by the Nevada Division of Public and Behavioral Health Bureau of Behavioral Health, Prevention, and Wellness through funding provided by the Nevada State Opioid Response Grant awarded by the Substance Abuse and Mental Health Services Administration.



What is an opioid and how can you overdose?

Opioids include illicit drugs such as heroin or prescription medications used to treat pain such as oxycodone, hydrocodone, and fentanyl. An opioid overdose can happen when you take too many opioids at once or mix them with other drugs and your breathing slows or stops.



To avoid an accidental opioid overdose:

- Avoid mixing your opioids with alcohol, benzodiazepines (Xanax, Ativan, Klonopin, Valium), or medicines that make you sleepy.
- Be extra careful if you miss or change doses, feel ill, or start new medications.



How to identify an opioid overdose:

Look for these common signs:

- The person won't wake up or respond to voice or touch
- Breathing slows or even stops
- Lips and fingernails turn blue or gray



In case of overdose:


1. Call 911 and administer naloxone.
2. Do rescue breathing or chest compressions
 - Check for clear airway. If blocked, roll on side and use finger sweep to clear.
 - Roll onto back, tilt head back and pinch nose.
 - Give 2 regular breaths.
 - If still not breathing on own, give 1 breath every 5 seconds.
 - Continue rescue breathing or chest compressions until person revives or help arrives.
 - Once the person starts breathing on own, put them in the recovery position. Remember to keep breathing for them until the person is able to breath on own.
3. Place into Recovery Position:
 - Roll the person on their side, with top arm placed under their head.
 - Pull the top knee and arm over their body to keep them on their side. This will help prevent them from choking if they vomit. After naloxone is administered, stay with a person for at least 3 hours or until help arrives.

How to administer injectable naloxone:

Step 1: Remove cap from naloxone vial and uncover the needle.

Step 2: Turn vial upside down and insert needle through rubber plug. Pull back on Plunger and draw 1ml (1cc) of liquid. Get rid of excess air.

Step 3: Inject into an upper arm, thigh muscle, or buttocks and empty the syringe.



Step 4: If no or minimal reaction in 2-3 minutes, give a second dose.

Nevada's Good Samaritan Law offers protection from arrest to individuals who seek medical assistance for a person experiencing a drug or alcohol overdose and the individual who experienced the overdose.

Protection is not offered from arrest for charges related to possession with intent to sell/distribute, child neglect/endangerment or for individuals with active warrants for their arrest.

**Now that you have naloxone . . .
Tell someone where it is and how to use it.**