

# How to test for Xylazine

## Powders & Pills guide

### Step 1: Prepare Sample

#### Crush Your Pill

- Make sure your pill or powder is crushed finely into a uniform consistency




### Step 2: Choose a Testing Method

#### Test Everything

- Pour powder into a cup
- Use 1/2 cup or 100ml of water per gram of material
- Mix until fully dissolved


#### Option 1 (Most Accurate)



#### Test Bag Residue

- Make sure the contents inside the bag are a uniform powder
- Dump the contents into another container
- Fill the empty bag with water and mix if necessary


#### Option 2



#### Test a Scoop

- Put one micro scoop of substance into a small cup
- Mix 1 tsp or 5ml of water for each scoop of substance


#### Option 3



## Step 3: Testing

### Use the Test Strip



- Place the test strip wavy side down in the water for 15s. Make sure the water is up to the line
- Remove strip from water and place it on a flat surface for 1-2 minutes



## Step 4: Results

### Read the Results

- A test **POSITIVE** for xylazine is indicated by **ONE RED LINE**
- A test **NEGATIVE** for xylazine is indicated by **TWO RED LINES**
- Remember, no test is 100% accurate, & a negative test does not necessarily mean you are safe



**NOTE:** It should be considered a **NEGATIVE** test even if a second red line is faded or faint  
If possible, use an additional test to confirm your results

